

A novel diagrammatic technique for the single-site Anderson model

V. A. Moskalenko^{1,2,*}, P. Entel³, D. F. Digor¹, L. A. Dohotaru⁴, and R. Citro⁵

¹*Institute of Applied Physics, Moldova Academy of Sciences, Chisinau 2028, Moldova*

²*BLTP, Joint Institute for Nuclear Research, 141980 Dubna, Russia*

³*University of Duisburg-Essen, 47048 Duisburg, Germany*

⁴*Technical University, Chisinau 2004, Moldova and*

⁵*Dipartimento di Fisica E. R. Caianiello, Università degli Studi di Salerno and CNISM, Unità di ricerca di Salerno, Via S. Allende, 84081 Baronissi (SA), Italy*

(Dated: February 6, 2008)

A diagrammatic theory around the atomic limit is proposed for the single-impurity Anderson model in which the strongly correlated impurity electrons hybridize with free (uncorrelated) conduction electrons. Using this diagrammatic approach, we prove the existence of a linked cluster theorem for the vacuum diagrams and derive Dyson type of equations for the localized and conduction electrons and corresponding equations for the mixed propagators. The system of equations can be closed by summing the infinite series of ladder diagrams containing irreducible Green's functions. The result allows to discuss resonances associated with the quantum transitions at the impurity site.

PACS numbers: 78.30.Am, 74.72.Dn, 75.30.Gw, 75.50.Ee

The study of strongly-correlated electron systems has become one of the most active fields of condensed matter physics during the last decade. The properties of these systems cannot be described by Fermi liquid theory. One of the important models of strongly correlated electrons is the single-site or impurity model introduced by Anderson in 1961 [1], which has been intensively discussed in many papers. It is a model for a system of free conduction electrons that interact with the system of a local spin of an electron in the d or f shells of an impurity atom. The impurity electrons are strongly correlated because of the strong on-site Coulomb repulsion and interact with the conduction electrons via exchange and hybridization. Most previous work is based on the method of equations of motion for retarded and advanced Green's functions and a truncating procedure as proposed, for example, by Bogoliubov and Tjablikov [2]. A first attempt to develop a diagrammatic theory for this problem was undertaken by Barabanov in [3]. With introduction of the dynamical mean field theory, the interest in the Anderson impurity model has considerably increased because infinite dimensional lattice models like the Hubbard model can be mapped onto effective impurity models and a self-consistency condition [4, 5].

The Hamiltonian of the model taking only s -like electrons into account is written as

$$\begin{aligned} H &= H_0 + H_{int}, \quad H_0 = H_0^c + H_0^f, \\ H_0^c &= \sum_{\mathbf{k}\sigma} \epsilon(\mathbf{k}) C_{\mathbf{k}\sigma}^+ C_{\mathbf{k}\sigma}, \quad H_0^f = \epsilon_f \sum_{\sigma} f_{\sigma}^+ f_{\sigma} + U n_{\uparrow}^f n_{\downarrow}^f, \\ H_{int} &= \frac{1}{\sqrt{N}} \sum_{\mathbf{k}\sigma} (V_{\mathbf{k}\sigma} f_{\sigma}^+ C_{\mathbf{k}\sigma} + V_{\mathbf{k}\sigma}^* C_{\mathbf{k}\sigma}^+ f_{\sigma}), \end{aligned} \quad (1)$$

where $n_{\sigma}^f = f_{\sigma}^+ f_{\sigma}$, $C_{\mathbf{k}\sigma}(C_{\mathbf{k}\sigma}^+)$ and $f_{\sigma}(f_{\sigma}^+)$ are the annihilation (creation) operators of conduction and impurity electrons with spin σ , respectively; $\epsilon(\mathbf{k})$ is the kinetic energy of the conduction band states $(\mathbf{k}, \sigma)$; ϵ_f is the lo-

cal energy of f -electrons and N is the number of lattice sites. H_{int} is the hybridization interaction between conduction and localized electrons. Summation over $\mathbf{k}$ will be changed to an integral over the energy $\epsilon(\mathbf{k})$ using the density of states $\rho_0(\epsilon)$ of conduction electrons.

The term in the Hamiltonian involving U describes the on-site Coulomb interaction between two impurity electrons. This term is far too large to be treated by perturbation theory. Therefore, we include it in the non-interacting Hamiltonian H_0 . Since this term invalidates Wick's theorem for local electrons, we first of all have to formulate the generalized Wick's theorem (GWT) for local electrons, preserving the ordinary Wick theorem for conduction electrons. Our GWT can be considered as the prescription which allows to determine the irreducible Green's functions or Kubo cumulants. A similar prescription has been used when discussing the properties of the single-band Hubbard model [6, 7, 8].

In interaction representation, the renormalized (Matsubara) Green's functions of conduction and impurity electrons have the form:

$$G(\mathbf{k}, \sigma, \tau | \mathbf{k}', \sigma', \tau') = -\langle T C_{\mathbf{k}\sigma}(\tau) \bar{C}_{\mathbf{k}'\sigma'}(\tau') U(\beta) \rangle_0^c, \quad (2)$$

$$g(\sigma, \tau | \sigma', \tau') = -\langle T f_{\sigma}(\tau) \bar{f}_{\sigma'}(\tau') U(\beta) \rangle_0^c. \quad (3)$$

Here τ and τ' stand for the imaginary time with $0 < \tau < \beta$ (β is the inverse temperature) and T is the chronological time ordering operator. The evolution operator $U(\beta)$ is determined by the hybridization interaction H_{int} . The statistical averaging in Eqs. (2) and (3) is carried out with respect to the zero-order density matrix of the conduction and impurity electrons.

The thermodynamic perturbation theory with respect to H_{int} requires an appropriate generalization scheme for the evaluation of statistical averages of the T -products of localized f -electron operators.

In zero-order approximation (neglecting the hy-

bridization term), the corresponding Green's functions have the form ($\omega \equiv \omega_n = (2n+1)\pi/\beta$):

$$G_{\sigma\sigma'}^0(\mathbf{k}, \mathbf{k}' | i\omega) = \delta_{\sigma\sigma'} \delta_{\mathbf{k}\mathbf{k}'} / \{i\omega - \epsilon(\mathbf{k})\},$$

$$g_{\sigma\sigma'}^0 = \delta_{\sigma\sigma'} g_{\sigma}^0(i\omega) = \delta_{\sigma\sigma'} \left\{ \frac{1 - n_{\bar{\sigma}}}{\lambda_{\sigma}(i\omega)} + \frac{n_{\bar{\sigma}}}{\bar{\lambda}_{\sigma}(i\omega)} \right\}, \quad (4)$$

where ($\bar{\sigma} = -\sigma$) and

$$\lambda_{\sigma}(i\omega) = i\omega + E_0 - E_{\sigma}, \quad \bar{\lambda}_{\bar{\sigma}}(i\omega) = i\omega + E_{\bar{\sigma}} - E_2,$$

$$Z_0 = e^{-\beta E_0} + e^{-\beta E_{\sigma}} + e^{-\beta E_{\bar{\sigma}}} + e^{-\beta E_2},$$

$$n_{\bar{\sigma}} = \{e^{-\beta E_{\bar{\sigma}}} + e^{-\beta E_2}\} / Z_0.$$

In the case of f -electrons, we can formulate the identity corresponding to the GWT:

$$\langle T f_1 f_2 \bar{f}_3 \bar{f}_4 \rangle_0 = \langle T f_1 \bar{f}_4 \rangle_0 \langle T f_2 \bar{f}_3 \rangle_0$$

$$- \langle T f_1 \bar{f}_3 \rangle_0 \langle T f_2 \bar{f}_4 \rangle_0 + \langle T f_1 f_2 \bar{f}_3 \bar{f}_4 \rangle_0^{ir}, \quad (5)$$

or

$$g_2^0(1, 2|3, 4) = g^0(1|4)g^0(2|3)$$

$$- g^0(1|3)g^0(2|4) + g_2^{(0)ir}(1, 2|3, 4), \quad (6)$$

where n stands for (σ_n, τ_n) . The generalization for more complicated averages of type $g_n^0(1, \dots, n | n+1, \dots, 2n) = (-1)^n \langle T f_1 \dots f_n \bar{f}_{n+1} \dots \bar{f}_{2n} \rangle_0$ is straightforward, i.e., the right-hand part of this relation will contain $n!$ terms of ordinary Wick type (chain diagrams) and also the different products of irreducible functions with the same total number of operators. These definitions are the simplification of the corresponding ones for Hubbard and other lattice models proposed earlier by us. The calculation of the simplest irreducible function like, for example, $g_2^{(0)ir}(12|34)$ is rather cumbersome but straightforward. Then, we determine the Fourier representation of $g_2^{(0)ir}[\sigma_1, \tau_1; \sigma_2, \tau_2 | \sigma_3, \tau_3; \sigma_4, \tau_4]$, for which exists spin and frequency conservation, $\sigma_1 + \sigma_2 = \sigma_3 + \sigma_4$, $\omega_1 + \omega_2 - \omega_3 - \omega_4 = 0$. Thus, we dispose of the rules to deal with chronological averages in thermodynamic perturbation theory.

First of all, we determine the thermodynamic potential F of the system,

$$F = F_0 - \frac{1}{\beta} \ln \langle U(\beta) \rangle_0, \quad (7)$$

where F_0 refers to the free impurity atom and conduction electrons. The diagrams which determine the thermodynamic potential have no external lines and represent the vacuum. Figure 1 shows the simplest connected vacuum diagrams of the normal (non-superconducting) state with the process of hybridization of C - and f -electrons. The zero order propagators of conduction and impurity electrons are represented by solid and dashed lines, respectively. These lines connect the crosses which depict the




FIG. 1: The simplest connected vacuum diagrams of the normal state. The diagram (a) is of second and (b) and (c) are of fourth order in the hybridization interaction. Here, $\bar{V}_n = V_n/\sqrt{N}$.

impurity states. To each of the crosses two arrows are attached, one of which is ingoing and the other outgoing. They stand for annihilation and creation of electrons. The index n denotes (σ_n, τ_n) for impurity and $(\mathbf{k}_n, \sigma_n, \tau_n)$ for conduction electrons. The rectangles with $2n$ crosses represent the irreducible $g_n^{(0)ir}$ Green's function, $g_n^{(0)ir}$.

Besides the vacuum diagrams of fourth order in Fig. 1 (b) and (c), there is also one disconnected diagram composed of two diagrams of type (a) containing an additional factor $1/2!$. This is repeated in high order of perturbation theory and permits us to formulate the linked cluster theorem, which has the form

$$\langle U(\beta) \rangle_0 = \exp \langle U(\beta) \rangle_0^c, \quad (8)$$

where $\langle U(\beta) \rangle_0^c$ contains only connected diagrams and is equal to zero when hybridization is switched off.

Now, we consider the diagrammatical analysis of the perturbation series for the renormalized propagator in Eq. (3). The simplest contributions to g in second order are represented in Fig. 2. All diagrams contain two external points with attached arrows fixed by the arguments of the Green's functions. At the inner points of the diagrams a summation over $\sigma_n, \mathbf{k}_n$ and integration over τ_n is included. The new diagrammatical elements which appear, are the irreducible two-particle Green's functions. The renormalization of these quantities is not considered here, only the renormalization of the propagators due to hybridization.




FIG. 2: The second order perturbation contribution to the impurity electron propagator.

The final equations for the renormalized Green's functions are more conveniently written down in Fourier rep-

resentation. For the propagator of conduction electrons we obtain:

$$G_{\sigma\sigma'}(\mathbf{k}, \mathbf{k}' | i\omega) = \delta_{\mathbf{k}\mathbf{k}'} \delta_{\sigma\sigma'} G_{\sigma}^0(\mathbf{k} | i\omega) + \frac{V_{\mathbf{k}}^* V_{\mathbf{k}'}}{N} G_{\sigma}^0(\mathbf{k} | i\omega) g_{\sigma\sigma'}(i\omega) G_{\sigma'}^0(\mathbf{k}' | i\omega). \quad (9)$$

This renormalized propagator is expressed by the full propagator g of the impurity electrons. Now, it is necessary to obtain the corresponding equation for the full impurity electron propagator. Because the subsystem of f -electrons is strongly correlated, we have to introduce the correlation function $Z_{\sigma\sigma'}$ which is represented by strongly connected diagrams containing irreducible Green's functions [6, 7, 8] and $\Lambda_{\sigma}(i\omega) = g_{\sigma}^0(i\omega) + Z_{\sigma}(i\omega)$. To this we introduce new quantities:

$$\frac{1}{N} \sum_{\mathbf{k}_1 \mathbf{k}_2} V_{\mathbf{k}_2}^* V_{\mathbf{k}_1} G^0(\mathbf{k}_1, \sigma_1, \tau_1 | \mathbf{k}_2, \sigma_2, \tau_2) = \frac{1}{N} \sum_{\mathbf{k}_1} |V_{\mathbf{k}_1}|^2 \times G_{\sigma_1\sigma_2}^0(\mathbf{k} | \tau_1 - \tau_2) \equiv \delta_{\sigma_1\sigma_2} G_{\sigma_1}^0(\tau_1 - \tau_2), \quad (10)$$

which allow to simplify the structure of equation for the f -electron propagator, which can be written as

$$g_{\sigma}(i\omega) = \{ \Lambda_{\sigma}(i\omega) - G_{\sigma}^0(-i\omega) \Lambda_{\sigma}(i\omega) \Lambda_{\sigma}(-i\omega) \} / d_{\sigma}(i\omega). \quad (11)$$

$$d_{\sigma}(i\omega) = [1 - \Lambda_{\sigma}(i\omega) G_{\sigma}^0(i\omega)] [1 - \Lambda_{\sigma}(-i\omega) G_{\sigma}^0(-i\omega)].$$

Equation (9) has originally been derived by Anderson [1] by using the equation of motion method in which the propagator $g_{\sigma}(i\omega)$ corresponds to the t -matrix for non-spin-flip scattering.

The main approximation which we employ here corresponds to the ladder approximation (which contain the main contributions of spin and charge fluctuations) and is schematically shown in Fig. 3. We investigate the prop-




FIG. 3: Diagrammatic representation of the approximative scheme used in the evaluation of the correlated function. The solid double lines with arrows depict the renormalized one-particle Green's function of conduction electrons. The rectangle depicts the irreducible Green's function of impurity electrons.

erties of the normal state of the single-site Anderson model by using Eqs. (9) and (11) for the corresponding Green's functions of conduction and impurity electrons. In the ladder approximation only the simplest irreducible Green's function $g_2^{(0)ir}$ is iterated many times. In the

paramagnetic phase ($\sigma' = \sigma$) the equation for Z reduces to

$$Z_{\sigma}(i\omega) = -\frac{1}{\beta} \sum_{\omega_1} \sum_{\sigma_1} G_{\sigma_1}(i\omega_1) \times \tilde{g}_2^{(0)ir}[\sigma, i\omega; \sigma_1, i\omega_1 | \sigma_1, i\omega_1; \sigma, i\omega], \quad (12)$$

where

$$G_{\sigma}(i\omega) = \frac{1}{N} \sum_{\mathbf{k}_1 \mathbf{k}_2} V_{\mathbf{k}_1}^* V_{\mathbf{k}_2} G_{\sigma}(\mathbf{k}_1, \mathbf{k}_2 | i\omega). \quad (13)$$

Using Eqs. (9) and (11) G can be written as

$$G_{\sigma}(i\omega) = G_{\sigma}^0(i\omega) + [G_{\sigma}^0(i\omega)]^2 g_{\sigma}(i\omega) = \frac{G_{\sigma}^0(i\omega)}{1 - \Lambda_{\sigma}(i\omega) G_{\sigma}^0(i\omega)}. \quad (14)$$

Then, using the definition of the correlation function, $\Lambda_{\sigma}(i\omega)$, and approximation (12) for $Z_{\sigma}(i\omega)$, we obtain the final integral equation for Λ_{σ} :

$$\Lambda_{\sigma}(i\omega) = g_{\sigma}^{(0)}(i\omega) - \frac{1}{\beta} \sum_{\omega_1} \sum_{\sigma_1} \frac{G_{\sigma_1}^0(i\omega_1)}{1 - \Lambda_{\sigma_1}(i\omega_1) G_{\sigma_1}^0(i\omega_1)} \times \tilde{g}_2^{(0)ir}[\sigma, i\omega; \sigma_1, i\omega_1 | \sigma_1, i\omega_1; \sigma, i\omega]. \quad (15)$$

For the symmetric case when $\epsilon_f = -U/2 < 0$ and $\epsilon_f + U = U/2 > 0$, we have more simple equations:

$$g_{\sigma}^0(i\omega) = i\omega \{ (i\omega)^2 - (U/2)^2 \},$$

$$Z_0 = 2(1 + \exp(\beta U/2)), \quad n_{\sigma} = 1/2 \quad (16)$$

together with the antisymmetry property of the zero-order impurity Green's function, $g_{\sigma}^0(-i\omega) = -g_{\sigma}^0(i\omega)$. In addition, if we assume that the matrix element $V(\epsilon)$ and the bare density of states, $\rho_0(\epsilon)$ are even function of its argument, then $G_{\sigma}^0(i\omega)$ is also antisymmetric, $G_{\sigma}^0(-i\omega) = -G_{\sigma}^0(i\omega)$. This allows us to search for the antisymmetric solution,

$$\Lambda_{\sigma}(-i\omega) = -\Lambda_{\sigma}(i\omega), \quad (17)$$

of Eq. (15). The analytical continuation of $G_{\sigma}^0(i\omega)$ has the form

$$G_{\sigma}^0(E + i\delta) = I(E) - i\Gamma(E),$$

$$I(E) = \text{P} \int d\epsilon \rho_0(\epsilon) \frac{|V(\epsilon)|^2}{E - \epsilon},$$

$$\Gamma(E) = \pi \rho_0(E) |V(E)|^2, \quad (18)$$

where $I(E)$ is the principal part of the integral. This function is antisymmetric. $\Gamma(E)$ is the band width of the virtual level and an even function of the energy. The symmetric impurity Anderson model is easier to treat

since it leads to a simple form for the irreducible two-particle Green's functions in the different spin and frequency channels yielding

$$\Lambda_\sigma(i\omega) = \frac{3(U/2)^2 G_\sigma^0(i\omega)}{[(i\omega)^2 - (U/2)^2]^2 [1 - \Lambda_\sigma(i\omega) G_\sigma^0(i\omega)]} + \frac{i\omega}{(i\omega)^2 - (U/2)^2}, \quad (19)$$

where we have used results obtained for the one-band Hubbard model [6, 7, 8]. There are two solutions of Eq. (19). The physical one with the correct asymptotic behavior, $\Lambda_\sigma(i\omega) \rightarrow 1/i\omega$ when $|\omega|$ tends to infinity, has the form:

$$\Lambda_\sigma(i\omega) = \frac{1}{2G_\sigma^0(i\omega)[(i\omega)^2 - (U/2)^2]} \times \left\{ [(i\omega)^2 - (U/2)^2 + i\omega G_\sigma^0(i\omega)] - [(i\omega)^2 - (U/2)^2 - i\omega G_\sigma^0(i\omega)] \sqrt{1 - 12Q(i\omega)} \right\}, \quad (20)$$

where

$$Q(i\omega) = \left(\frac{(U/2)G_\sigma^0(i\omega)}{(i\omega)^2 - (U/2)^2 - i\omega G_\sigma^0(i\omega)} \right)^2. \quad (21)$$

Finally, we obtain for the renormalized impurity electron propagator:

$$g(i\omega) = 2 \left\{ g^0(i\omega)/G^0(i\omega) + 3Q(i\omega)[g^0(i\omega) - 1/G^0(i\omega)]^2 \right\} \left\{ [1/G^0(i\omega) - g^0(i\omega)][1 + \sqrt{1 - 12Q(i\omega)}] + 6Q(i\omega)\{g^0(i\omega)G^0(i\omega) - 1\} \right\}^{-1}. \quad (22)$$

From the analytical continuation we obtain the spectral function of impurity electrons, which is given in Hubbard I approximation by

$$A_f^I(E) = \frac{2\Gamma(E)}{[E - (U/2)^2/E - I(E)]^2 + \Gamma^2(E)}. \quad (23)$$

Equation (23) has two resonances at $E = \pm E_0$, where E_0 is determined from the condition that the bracket in the denominator of (23) is equal to zero. These resonances are broadened by $\Gamma(E)$ which determines height and width of both resonances. This approximation yields a vanishing spectral function on the Fermi surface, where $E = 0$. Only beyond Hubbard I approximation, when strong correlations are taken into account, we obtain from Eq. (20) the spectral function of impurity electrons in the vicinity of the Fermi level having the following form,

$$A_f(E) \simeq \frac{12(2\Gamma/U)^2/\Gamma}{1 + 6(2\Gamma/U)^2 + \sqrt{1 + 12(2\Gamma/U)^2}}. \quad (24)$$




FIG. 4: Spectral function $A_f(E) \times W$ for different values of the theory parameters as function of energy E/W in the Hubbard I approximation (case (a)) and in our ladder approximation (cases (b) and (c)). In the cases (a) and (b) $U = 1 \text{ eV}$, $W = 2 \text{ eV}$, $\Gamma = 0.1 \text{ eV}$ and in case case (c) $U = 1.5 \text{ eV}$, $W = 2 \text{ eV}$, $\Gamma = 0.15 \text{ eV}$.

In conclusion we state that our diagrammatic approach allows to describe in ladder approximation for the irreducible Green's function two resonances at energies $E = \pm E_0$ and the feature at $E = 0$ given by Eq. (24). The distance between two peaks (Figure 4) is determined by the value of U but the height and the width of the peaks by Γ . Beyond the Hubbard I approximation the non zero values of spectral function appear for $E = 0$.

This work has been supported by a Grant from the Heisenberg-Landau Program. V. A. M. would like to thank the Universities of Duisburg-Essen and Salerno for hospitality and financial support.

* Electronic address: moskalen@thsun1.jinr.ru

- [1] P. W. Anderson, Phys. Rev. **124**, 41 (1961).
- [2] N. N. Bogoliubov and S. V. Tjablikov, Doklady AN USSR, **126**, 53 (1959) [in Russian].
- [3] A. F. Barabanov, C. A. Kikoin and L. A. Maximov, Theor. Math. Phys. **20**, 364 (1974).
- [4] A. Georges, G. Kotliar, W. Krauth and M. J. Rozenberg, Rev. Mod. Phys. **8**, 13 (1996).
- [5] G. Kotliar and D. Vollhardt, Physics Today **57**, 53 (2004).
- [6] M. I. Vladimir and V. A. Moskalenko, Theor. Math. Phys. **82**, 301 (1990).
- [7] S. I. Vakar, M. I. Vladimir and V. A. Moskalenko, Theor. Math. Phys. **85**, 1185 (1990).
- [8] V. A. Moskalenko, P. Entel and D. F. Digor, Phys. Rev. B **59**, 619 (1999).